



PubMed Nucleotide Protein Genome Structure PopSet Taxonomy OMIM Bo

Search PubMed for

Limits

Preview/Index

History

Clipboard

Details

About Entrez

Display

Abstract

Sort

Save

Clip Add

Order

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

PubMed Services

Journal Browser

MeSH Browser

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

☐ 1: Arthritis Rheum 2001 Dec;44(12):2836-40Related Articles, **NEW Books**,
LinkOut**Response of Wegener's granulomatosis to anti-CD20 chimeric monoclonal antibody therapy.****Specks U, Fervenza FC, McDonald TJ, Hogan MC.**

Division of Pulmonary and Critical Care Medicine, Mayo Clinic and Foundation, Rochester, MN 55905, USA. specks.ulrich@mayo.edu

We report on the successful, compassionate use of the anti-CD20 chimeric monoclonal antibody rituximab in a patient with chronic, relapsing cytoplasmic antineutrophil cytoplasmic antibody (cANCA)-associated Wegener's granulomatosis (WG). The patient initially responded to treatment with glucocorticoids and cyclophosphamide. However, bone marrow toxicity during cyclophosphamide treatment of a relapse precluded its further use. Azathioprine and mycophenolate mofetil treatment had failed to maintain remission of the WG, and methotrexate was contraindicated. Because the patient's 5-year course was characterized by close correlation of cANCA levels with disease activity, selective elimination of cANCA was deemed a treatment option for his latest relapse. He was given 4 infusions of 375 mg/M2 of rituximab and high-dose glucocorticoids. Complete remission was associated with the disappearance of B lymphocytes and cANCA. Glucocorticoid treatment was then discontinued. After 11 months, the cANCA recurred, and rituximab therapy was repeated, without glucocorticoids. At 8 months after the second course of rituximab (18 months after the first course), the patient's WG has remained in complete remission. Elimination of B cells by rituximab therapy may prove to be an effective and safe new treatment modality for ANCA-associated vasculitis and possibly other autoimmune diseases. This modality warrants closer examination in a carefully conducted clinical trial.

PMID: 11762944 [PubMed - indexed for MEDLINE]

Display

Abstract

Sort

Save

Text

Clip Add

Order